

## Definitions and sources (abbreviations given in footnote)

Group	Indicator	Description	Numerator	Denominator	Source
Deprivation	<b>South African Index of Multiple Deprivation Rank (1 = most deprived)</b>	Ranking for composite index of deprivation. Indicators cover the following domains: income and material deprivation, employment deprivation, education deprivation, and living environment deprivation	Antenatal 1st visit - total	Estimated pregnant women ~10 weeks' gestation	Southern African Social Policy Research Institute (SASPRI), based on data from Stats SA Census 2011
	<b>Antenatal 1st visit coverage</b>	The proportion of potential antenatal clients coming for at least one (booking) antenatal visit. The census number of children under one year factorised by 1.15 is used as a proxy denominator – the extra 0.15 (15%) is a rough estimate to cater for late miscarriages (~10 to 26 w), stillbirths (after 26 weeks' gestation) and infant mortality. Pregnant women are regarded as potential antenatal clients from around 10 weeks' gestation, i.e. spontaneous abortions before that as well as TOP cases are excluded			webDHIS
	<b>Delivery in 10–19 years in facility rate</b>	Deliveries where the mother is 10–19 years old and such delivery is conducted by a trained healthcare worker in a health facility	Deliveries among women aged 10–19 years	Number of deliveries - total	webDHIS
	<b>Maternal mortality in facility ratio</b>	Women who died in hospital as a result of childbearing, during pregnancy or within 42 days of delivery or termination of pregnancy, per 100 000 live births in facility	Maternal deaths in facility	Live births in facility	webDHIS
	<b>Early neonatal death in facility rate</b>	Early neonatal deaths per 1000 infants who were born alive in health facilities.	Death in facility 0–6 days	Live births in facility	webDHIS
	<b>Neonatal death in facility rate</b>	Infants 0–28 days who died during their stay in the facility as a proportion of live births in facility	Neonatal 0–28 days death in facility	Live births in facility	webDHIS
	<b>Child under 5 years pneumonia case fatality rate</b>	Proportion of children under 5 years admitted with pneumonia who died	Child under 5 years pneumonia death	Child under 5 years pneumonia admitted	webDHIS
	<b>Death under 5 years against live birth rate</b>	Children under 5 years who died during their stay in the facility as a proportion of all live births	Death in facility under 5 years total	Live births in facility	webDHIS
	<b>Immunisation coverage under 1 year</b>	Percentage of all children in the target area under 1 year who complete their primary course of immunisation. A primary course comprises BCG, OPV 1, DTaP-IVP-Hib-HBV 1, 2 and 3, PCV 1,2 and 3, RV 1 and 2 and measles 1	Children fully immunised under 1 year	Target population under 1 year	webDHIS
	<b>Measles 2nd dose coverage</b>	Children 1 year (12 months) who received measles 2nd dose, as a proportion of the 1 year population.	Measles 2nd dose	Target population 1 year	webDHIS

Group	Indicator	Description	Numerator	Denominator	Source
Women's health	Couple year protection rate	Women protected against pregnancy by using modern contraceptive methods, including sterilisations, as proportion of female population 15–49 year. Contraceptive years are the total of (Oral pill cycles / 15) + (Medroxyprogesterone injection / 4) + (Norethisterone enanthate injection / 6) + (IUCD x 4.5) + (Sub-dermal implant x 2.5) + Male condoms distributed / 120) + (Female condoms distributed / 120) + (Male sterilisation x 10) + (Female sterilisation x 10)	Contraceptive years dispensed (including sterilisations)	Female target population 15–44 years	webDHIS
	TB symptom child under 5 years screened in facility rate	Children under 5 years screened for TB symptoms as a proportion of PHC headcount under 5 years	Child under 5 years screened for TB symptoms	PHC headcount under 5 years	webDHIS
	TB symptom 5 years and older screened in facility rate	Clients 5 years and older screened for TB symptoms as a proportion of PHC headcount 5 years and older	Client 5 years and older screened for TB symptoms	PHC headcount 5 years and older	webDHIS
	TB drug-susceptible successful treatment rate	Proportion TB clients (ALL types of TB) cured plus those who completed treatment	TB client cured OR completed treatment	TB client initiated on treatment	webDHIS
	TB drug-susceptible client loss to follow-up rate	TB clients who started drug-susceptible tuberculosis treatment and who subsequently became loss to follow-up as a proportion of all those in the treatment outcome cohort	All drug-susceptible TB loss to follow-up	All drug-susceptible TB patients in treatment outcome cohort	webDHIS
	TB drug-susceptible death rate	TB clients who started drug-susceptible tuberculosis treatment and who subsequently died as a proportion of all those in the treatment outcome cohort	All drug-susceptible TB client died	All drug-susceptible TB patients in treatment outcome cohort	webDHIS
	TB multidrug-resistant treatment success rate	TB-MDR client successfully completing treatment as a proportion of TB-MDR confirmed clients started on treatment	TB-MDR client successfully complete treatment	TB-MDR confirmed client start on treatment	webDHIS
	TB-XDR treatment success rate	TB-XDR clients successfully completing treatment as a proportion of TB-XDR confirmed clients started on treatment	TB-XDR client successfully complete treatment	TB-XDR confirmed client start on treatment	webDHIS
	Clients remaining on ART rate	Percentage of estimated people living with HIV who remain on ART	Total clients remaining on ART at end of the month	Estimated number of people living with HIV	webDHIS
HIV	Adult with viral load suppressed rate 12 months	Proportion of adult ART clients with viral load suppressed at 12 months	ART client viral load under 400 cps/mL at 12 months	ART viral load done	webDHIS
	Child with viral load suppressed rate 12 months	Proportion of ART clients with viral load suppressed at different time intervals. This indicates the population level immunological impact of clients on ART.	ART Child Viral Load Suppressed (VLS)	ART Child client viral load done (VLD)	webDHIS
	ART client viral load suppressed rate (VLS)	ART viral load suppressed - total as a proportion of ART viral load done - total	ART viral load suppressed - total	ART viral load done - total	webDHIS

Group	Indicator	Description	Numerator	Denominator	Source
Non-communicable diseases	Cervical cancer screening coverage	Cervical smears in women 30 years and older as a proportion of the female population 30 years and older. 80% of these women should be screened for cervical cancer every 10 years and 20% must be screened every 3 years	Cervical cancer screening in non- HIV woman 30 years and older + cervical cancer screening in women 20 years and older	80% women aged 30-59yrs / 10 + 20% women aged 20-59yrs / 3	webDHIS
	PHC utilisation rate	The rate at which PHC services are utilised by clients of all ages in the catchment population, represented as the average number of PHC visits per client per year in the target population. The denominator is usually Census-derived population	Total PHC headcount	Total population	webDHIS
Inpatient management	Average length of stay	Average number of patient days that an admitted patient spends in hospital before separation	Inpatient days + 1/2 day patients	Separations = Discharges (including day patients) + Deaths + Transfers out	webDHIS
	Inpatient bed utilisation rate	Inpatient bed days used as proportion of maximum Inpatient bed days (inpatient beds x days in period) available	Total patient days = (Inpatient days + 1/2 day patients) x 100	Total usable bed days = (Inpatient beds – total) x 30.42	webDHIS
	Patient day equivalent	Weighted data element as proxy for estimating resources for all types of patients in terms of inpatient days	The sum of inpatient days total x 1, day patient total x 0.5, and OPD/emergency total headcount x 0.3333333	-	webDHIS
	Inpatient crude death rate	Proportion of admitted patients/separations who died during hospital stay	Inpatient deaths – total	Separations = Discharges (including day patients) + Deaths + Transfers out	webDHIS
	Hospital beds per 10 000 target population	Number of inpatient beds per 10 000 target population. For public sector beds, the uninsured population is used as the target	Inpatient beds	Uninsured population (total population less medical scheme coverage x population)	webDHIS

Group	Indicator			Description			Numerator			Denominator			Source			
	Provincial and Local Government expenditure on District Health Services per capita (uninsured)	Provincial PHC expenditure per capita (uninsured)	Provincial expenditure per PHC headcount	Total amount spent on DHS per person without medical scheme coverage	Total amount spent on non-hospital PHC health services per person without medical scheme coverage	Total amount spent on non-hospital PHC health services per headcount	Provincial expenditure on DHS (excluding 2.8 Coroner) plus net LG expenditure on PHC	Provincial expenditure on PR2.2-2.7 of DHS + net LG expenditure	Provincial expenditure on PR2.2-2.7 of DHS + net LG expenditure	Uninsured population (total population less medical scheme coverage x population)	Uninsured population (total population less medical scheme coverage x population)	Total PHC headcount	Uninsured population (total population less medical scheme coverage x population)	Uninsured population (total population less medical scheme coverage x population)	Uninsured population (total population less medical scheme coverage x population)	Uninsured population (total population less medical scheme coverage x population)
Finance																Calculated from BAS, Treasury data on LG expenditure, webDHS population and GHS modelled for medical scheme coverage
																Calculated from BAS, Treasury data on LG expenditure, webDHS population and GHS modelled for medical scheme coverage
																BAS, webDHS
Human resources																BAS, webDHS
																Calculated from PERSAL, webDHS population and GHS modelled estimates of medical scheme coverage
																Calculated from PERSAL, webDHS population and GHS modelled estimates of medical scheme coverage
																Calculated from PERSAL, webDHS population and GHS modelled estimates of medical scheme coverage
Extra																GHS modelled estimates of medical scheme coverage

ART = antiretroviral treatment; BAS = National Treasury Basic Accounting System; BCG = bacille Calmette-Guerin; BP = blood pressure; CDC = Community Day Centre; CHC = Community Health Centre; DHIS = District Health Information System; DHS = District Health Services; DTaP-IVP-Hib-HBV = Diphtheria, Tetanus, Inactivated Polio Vaccine, Haemophilus Influenza B, Hepatitis Vaccine; GHS = General Household Survey; HIV = Human Immunodeficiency Virus; LG = Local Government; MDR = multidrug-resistant; NCDs = non-communicable diseases; NDoH = National Department of Health; OPD = outpatient department; OPV = oral polio vaccine; PCR = polymerase chain reaction; PCV = pneumococcal conjugate vaccine; PDE = patient day equivalent; PHC = Primary Health Care; PR = programme; RV = Rotavirus vaccine;; Stats SA = Statistics South Africa; TB = tuberculosis; ToP = termination of pregnancy; XDR = extremely drug-resistant;